




Central Diffractive Processes
at the Tevaron and LHC.



V.A. Khoze (IPPP, Durham & PNPI)



(In collaboration with L. Harland-Lang, M. Ryskin and W.J. Stirling)



χ_c, χ_b




- Introduction. _____
- Central exclusive production (CEP) of $\chi_{c0,1,2}$ states at the Tevatron, LHC and RHIC.
- Overview of $\gamma\gamma$ and χ_b CEP results and ongoing studies.
- Forward proton distributions and correlations.
- CDP@LHCb with FSC ?
- Conclusion.

Introduction

Why are we interested in central exclusive χ_c ($\chi_b, \gamma\gamma, jj$) production?

- Driven by same mechanism as Higgs (or other new object) CEP at the LHC.

- χ_c, jj and $\gamma\gamma$ CEP has been observed by CDF. 

New D0 results.
RHIC data
to come (hopefully) soon

→ Can serve as ‘Standard Candle’ processes, which allow us to check the theoretical predictions for central exclusive new physics signals at the LHC.

- $\chi_{c,b}$ production is of special interest: (star reactions!)
 - Heavy quarkonium production can shed light on the physics of bound states (lattice, NRQCD...).
 - Potential to produce different J^P states, which exhibit characteristic features (e.g. angular distributions of forward protons).
 - Could perhaps shed light on the various ‘exotic’ charmonium states observed recently.

Spin-Parity Analyzer




Detailed tests of dynamics of soft diffraction (KMR-02)



“soft” scattering can easily destroy the gaps

$S^2 \rightarrow$ absorption effects -necessitated by unitarity

Everybody's ~ happy (KMR, GLMM, FHSW, KP, Petrov et al, BH, GGPS, Luna...MCs)



eikonal rescatt: between protons
 enhanced rescatt: involving intermediate partons

soft-hard factorizⁿ
 conserved
 broken


Subject of hot discussions nowadays : S^2_{enh}



Standard Candle Processes

'BETTER TO LIGHT A CANDLE THAN TO
RANT AGAINST DARKNESS'

(Confucius)



The process $p-p \rightarrow \gamma\gamma / \chi_c / \chi_b / jj$ are standard candles for the exclusive Higgs

像教行子孔師先




孔夫子

孔丘 Kong Qiu

Bottomonium history started 30 years ago

(PRL 39, 242 (1977) and PRL 39,1240 (1977))

30 years later....




$M(\Upsilon) = 9.40 \pm 0.013$

$M(\Upsilon') = 10.00 \pm 0.04$

$M(\Upsilon'') = 10.43 \pm 0.12$

FNAL, E288

(spins- still unconfirmed)



$Y(nS)$ confirmed and $\chi_{bJ}(1,2P)$ states observed

Below $B\bar{B}$, several states not yet observed: 3 S-wave (η_b), 2 P-wave (h_b), D-wave, etc...

(Currently no complete theoretical description of onium properties.)

$\eta_b(1S)$

(BABAR (2008))

(Still puzzles)



The heaviest and most compact quark-antiquark bound state in nature



(Cannot detect p/pbar, down beam pipe, but BSC → η = 7.4 empty)

FSC@LHC

$$p + \bar{p} \rightarrow p + \gamma + \bar{p}$$

Cleanest (no S.I.) but smallest σ

KMR: 38 pb in our box). 2+1 candidates (more coming soon)

$$p + \bar{p} \rightarrow p + \chi_c + \bar{p}$$

Clean, big σ:

$$\frac{d\sigma}{dy}(y=0) \sim 100 \text{ nb (KMRS)}$$

$$p + \bar{p} \rightarrow p + \chi_b + \bar{p}$$

but M(c) small (non-pert) & hadron

$$p + \bar{p} \rightarrow p + JJ + \bar{p}$$

More perturbative, smaller theory uncertainty
But σ ~ 1/500th χ_c. Also BR's not known!

Prprospects !


Big cross section, but least well defined (jets!)
and largest background. ~ 100 pb for M(JJ) > 30 GeV

Our 3 measurements are all in good agreement
(factor “few”) with the Durham group predictions.

We set out to measure exclusive $\chi_{c0} \rightarrow J/\psi + \gamma \rightarrow \mu^+ \mu^- \gamma$

BSC very important as rap gap detectors. All LHC experiments should have them!

(Gap Detectors in no P-U events)



& nothing else
in all CDF
 $-7.4 < |\eta| < +7.4$

Beam Shower Counters BSC: $5.2 < |\eta| < 7.4$

If these are all empty, p and \bar{p} did not dissociate

but went down beam pipe with small ($\lesssim 1$ GeV/c) transverse momentum.



Mike Albrow

Exclusive production in CDF


Diffraction Trento Jan 2010

Forward physics with rapidity gaps at the LHC

M.G. Albrow,^{a,1} A. De Roeck,^b V.A. Khoze,^c J. Lämsä,^{d,e} E. Norbeck,^f Y. Onel,^f
R. Orava,^e A. Penzo^e and M.G. Ryskin^b




Accessible warm beam pipe
between BMX magnets



Can put scintillators at several z-locations
FSC = Forward Shower Counters

Do not see primary particles, but showers in pipe and other material.

Example, O_{++} -case



$$T = A\pi^2 \int \frac{d^2Q_{\perp} P(\chi(0^+))}{Q_{\perp}^2 (\vec{Q}_{\perp} - \vec{p}_{1\perp})^2 (\vec{Q}_{\perp} + \vec{p}_{2\perp})^2} f_g(x_1, x'_1, Q_1^2, \mu^2; t_1) f_g(x_2, x'_2, Q_2^2, \mu^2; t_2),$$

$$A^2 = 8\pi\Gamma(\chi \rightarrow gg)/M_{\chi}^3 \quad *K_{\text{NLO}} \quad P(\chi(0^+)) = (\vec{Q}_{\perp} - \vec{p}_{1\perp}) \cdot (\vec{Q}_{\perp} + \vec{p}_{2\perp}).$$

● Strong sensitivity to the polarization structure of the vertex in the bare amplitude.

KMR-01

Absorption is sizeably distorted by the polarization structure (affects the b-space distr.)

● χ_c, χ_b -production is especially sensitive to the effects of enhanced absorption

■ larger available rapidity interval

■ lower scale \rightarrow larger dipole size \rightarrow larger absorption

(Gap size for χ_c at the Tevatron is expected to exceed that for the Higgs at the LHC)

KMR-02, KKMR-03,
HKRS 09-10

● Forward proton distributions & correlations- possibility to test diffraction dynamics

KMR-02


- 65 ± 10 signal χ_c events observed, but with a limited $M(J/\psi\gamma)$ resolution.
- Possible contribution from χ_{c1} and χ_{c2} states assumed, rather than observed, to be negligible.
- Assuming χ_{c0} dominance, CDF found:

$$\left. \frac{d\sigma(\chi_{c0})}{dy_\chi} \right|_{y=0} = (76 \pm 14) \text{ nb} ,$$

in good agreement with the previous KMRS value of 90 nb
([arXiv:0403218](https://arxiv.org/abs/0403218)).

Too good to be true ?!

- But can we be sure that χ_{c1} and χ_{c2} events do not contribute?





- A new MC (available on HepForge) including:
 - Non-forward $p_{\perp} \neq 0$ protons via the 'effective' slope parameters b_{eff} .
 - Full simulation of $\chi_{c(0,1,2)}$ CEP via the $\chi_c \rightarrow J/\psi\gamma \rightarrow \mu^+\mu^-\gamma$ decay chain.
 - $\chi_{b(0,1,2)}$ CEP via the equivalent $\chi_b \rightarrow \Upsilon\gamma \rightarrow \mu^+\mu^-\gamma$ decay chain.
 - More to come...
- The angular distributions of the final state particles, modeled in the MC, might help us to distinguish between the different states...
- ...however the severity of current CDF experimental cuts for χ_c CEP ($p_{\perp}(\mu) > 1.4 \text{ GeV}/c$, $|\eta_{\mu}| < 1$) appears to preclude this.
- **Diphoton CEP**


χ_{c1} and χ_{c2} : general considerations

- General considerations tell us that χ_{c1} and χ_{c2} CEP rates are strongly suppressed:
 - χ_{c1} : Landau-Yang theorem forbids decay of a $J = 1$ particle into on-shell gluons.
 - χ_{c2} : Forbidden (in the non-relativistic quarkonium approximation) by $J_z = 0$ selection rule that operates for forward ($p_{\perp} = 0$) outgoing photons. KMR-01 (A. Alekseev-1958-positronium)
- However the experimentally observed decay chain $\chi_c \rightarrow J/\psi \gamma \rightarrow \mu^+ \mu^- \gamma$ strongly favours $\chi_{c(1,2)}$ production, with:

$$\text{Br}(\chi_{c0} \rightarrow J/\psi \gamma) = 1.1\% ,$$

$$\text{Br}(\chi_{c1} \rightarrow J/\psi \gamma) = 34\% ,$$

$$\text{Br}(\chi_{c2} \rightarrow J/\psi \gamma) = 19\% .$$

- We should therefore seriously consider the possibility of $\chi_{c(1,2)}$ 

(R.Pasechnik et al, Phys.Lett.B680:62-71,2009; HKRS, Eur.Phys.J.C65:433-448,2010)

Cross section results (1)

- We find the following approximate hierarchy for the spin-summed amplitudes squared (assuming an exponential proton form factor $e^{-b\mathbf{p}_\perp^2}$).

$$|V_0|^2 : |V_1|^2 : |V_2|^2 \sim 1 : \frac{\langle \mathbf{p}_\perp^2 \rangle}{M_\chi^2} : \frac{\langle \mathbf{p}_\perp^2 \rangle^2}{\langle \mathbf{Q}_\perp^2 \rangle^2}. \quad (2)$$

- This $\sim 1/40$ suppression for the $\chi_{c1,2}$ states will be compensated by the larger $\chi_c \rightarrow J/\psi\gamma$ branching ratios, as well as by the larger survival factors S_{eik}^2 for the more peripheral reactions.
- An explicit calculation gives (for the perturbative contribution):

$$\frac{\Gamma_{J/\psi+\gamma}^{\chi_0}}{\Gamma_{\text{tot}}^{\chi_0}} \frac{d\sigma_{\chi_{c0}}^{\text{pert}}}{dy} : \frac{\Gamma_{J/\psi+\gamma}^{\chi_1}}{\Gamma_{\text{tot}}^{\chi_1}} \frac{d\sigma_{\chi_{c1}}^{\text{pert}}}{dy} : \frac{\Gamma_{J/\psi+\gamma}^{\chi_2}}{\Gamma_{\text{tot}}^{\chi_2}} \frac{d\sigma_{\chi_{c2}}^{\text{pert}}}{dy} \approx 1 : 0.6 : 0.22$$

- Note: these approximate values carry a factor of $\sim \frac{\times}{\div} 2$ uncertainty.



Cross section results

- As the cms energy increases we have:
 - Larger gluon density at smaller x values.
 - Smaller S_{eik}^2 survival factor.
 - Smaller S_{enh}^2 due to increase in size of rapidity gaps ($\sim s/m_\chi^2$) available for 'enhanced' absorptive effects.
- The combined result of these different effects is that the χ_c CEP rate has only a very weak energy dependence going from the Tevatron to the LHC.

• S_{eik}^2 and S_{enh}^2 accounted for in the integrand

\sqrt{s} (TeV)	0.5	1.96	7	10	14
$\frac{d\sigma}{dy_{\chi_c}}(pp \rightarrow pp(J/\psi + \gamma))$	0.57	0.73	0.89	0.92	1.0
$\frac{d\sigma(1^+)}{d\sigma(0^+)}$	0.59	0.61	0.69	0.67	0.71
$\frac{d\sigma(2^+)}{d\sigma(0^+)}$	0.21	0.22	0.23	0.23	0.23

Differential cross section (in nb) at rapidity $y_\chi = 0$ for central exclusive χ_{cJ} production via the $\chi_{cJ} \rightarrow J/\psi\gamma$ decay chain, summed over the $J = 0, 1, 2$ contributions, at RHIC, Tevatron and LHC energies, and calculated using GRV94HO partons,

$\chi_c \rightarrow \pi\pi, \chi_c \rightarrow K\bar{K}$

Spin-parity Analyzer

$$\text{BR}(\chi_{b1} \rightarrow \Upsilon\gamma) = (35 \pm 8)\%$$

$$\text{BR}(\chi_{b2} \rightarrow \Upsilon\gamma) = (22 \pm 4)\%$$

- Calculation exactly analagous to χ_c case with same hierarchy :
 However we have a stronger supression in the χ_{b1} and χ_{b2} rates than for the χ_c case.

- Larger $\langle Q_{\perp}^2 \rangle$ scale gives smaller b_{eff} values, i.e. non-forward effects are less strong, but still important.

- Significant uncertainties in input parameters:

- Only have $\text{Br}(\chi_{b0} \rightarrow \Upsilon\gamma) < 6\%$ from experiment (Crystal Ball -1986)
- $\Gamma_{\text{tot}}(\chi_{b0})$ experimentally undetermined. 🤔

- Consistently with the results of NRQCD, as well as the existing experimental data, we can take the values³ $\Gamma(\chi_{b0} \rightarrow gg) = 0.8$ MeV and $\text{Br}(\chi_{b0} \rightarrow \Upsilon\gamma) = 3\%$.

- $\chi_b(nP) \rightarrow DX$ (about 0.25 of all hadronic decays (CLEO-2009)

$$\chi_{b1} \rightarrow c\bar{c}X \quad (\text{Barbieri et al (1979), NRQCD})$$

FSC@LHCb ?



Suppressed non-resonant background $\sim m_c^2/M_{\chi_b}^2$

χ_b CEP (2)

$$\frac{\Gamma_{\Upsilon+\gamma}^{\chi_0} \frac{d\sigma_{\chi_{b0}}^{\text{pert}}}{dy}}{\Gamma_{\text{tot}}^{\chi_0}} : \frac{\Gamma_{\Upsilon+\gamma}^{\chi_1} \frac{d\sigma_{\chi_{b1}}^{\text{pert}}}{dy}}{\Gamma_{\text{tot}}^{\chi_1}} : \frac{\Gamma_{\Upsilon+\gamma}^{\chi_2} \frac{d\sigma_{\chi_{b2}}^{\text{pert}}}{dy}}{\Gamma_{\text{tot}}^{\chi_2}} \approx \mathbf{1 : 0.03 : 0.08}$$

\sqrt{s} (TeV)	0.5	1.96	7	10	14
$\frac{d\sigma}{dy_{\chi}}(\chi_{c0})$	27	35	42	43	45
$\frac{d\sigma}{dy_{\chi}}(\chi_{b0})$	-	0.017	0.021	0.022	0.022

Table 4: Differential cross section (in nb) at rapidity $y_{\chi} = 0$ for central exclusive $\chi_{(b,c)0}$ production at RHIC, Tevatron and LHC energies, and calculated using GRV94HO partons, as explained in the text.

\sqrt{s} (TeV)	1.96	7	10	14
$\frac{d\sigma}{dy_{\chi_b}}(pp \rightarrow pp(\Upsilon + \gamma))$	0.56	0.70	0.73	0.74
$\frac{d\sigma(1^+)}{d\sigma(0^+)}$	0.029	0.032	0.032	0.034
$\frac{d\sigma(2^+)}{d\sigma(0^+)}$	0.077	0.081	0.081	0.083

Table 5: Differential cross section (in pb) at rapidity $y_{\chi} = 0$ for central exclusive χ_{bJ} production via the $\chi_{bJ} \rightarrow \Upsilon\gamma$ decay chain, summed over the $J = 0, 1, 2$ contributions, at Tevatron and LHC energies, and calculated using GRV94HO partons, as explained in the text.

Measuring forward proton angular distributions

KKMR-03

- For low proton transverse momenta $p_{1,2\perp}$ we have:

$$p_{\perp}^2 \ll Q^2$$

$$d\sigma(0^+)/d\phi \approx \text{const.},$$


$$d\sigma(1^+)/d\phi \approx (p_{1\perp} - p_{2\perp})^2,$$

$$d\sigma(0^-)/d\phi \approx p_{1\perp}^2 p_{2\perp}^2 \sin^2 \phi,$$

while there does not exist a simple closed form for the χ_2 case

- Note these will receive corrections of $O(p_{\perp}^2 / \langle Q_{\perp}^2 \rangle)$.
- These distributions are strongly affected by absorptive corrections, through their dependence on the proton distribution in impact parameter b space.
- Forward proton detection would allow a clear discrimination between the different J states.

Very topical for STAR@RHIC forthcoming measurements with tagged forward protons
(new HKRS results soon to come).



Distribution (in arbitrary units) within the perturbative framework of the outgoing proton $p_{1\perp}^2$, integrated over the second proton $p_{2\perp}$, for the CEP of different J^P $\mathcal{C}\mathcal{P}$ states at $\sqrt{s} = 14$ TeV. The solid (dotted) line shows the distribution including (excluding) the survival factor, calculated using the two channel eikonal model of Ref. [74], while the dashed line shows the distribution in the small p_{\perp} limit, using the vertices of Eqs. (3.16)–(3.18) and excluding the survival factor.





Figure 4: Distribution (in arbitrary units) within the perturbative framework of the difference in azimuthal angle of the outgoing protons for the CEP of different J^P $c\bar{c}$ states at $\sqrt{s} = 14$ TeV. The solid (dotted) line shows the distribution including (excluding) the survival factor, calculated using the two channel eikonal model of Ref. [74], while the dashed line shows the distribution in the small p_{\perp} limit, using the vertices of Eqs. (3.16)–(3.18) and excluding the survival factor.

(KMRS, [arXiv:0409037](#))

- 3 candidate events observed by CDF ([arXiv:0707.237](#)), with more to come.
- More events would allow us to probe scaling of σ with E_{cut} .
- Similar uncertainties to χ_c case for low E_{cut} scale.
- Potential $|J_Z| = 2$ contribution found to be unimportant.
- New encouraging results for $gg \rightarrow \pi^0 \pi^0$ background.
- $\gamma\gamma$ CEP now included in SuperCHIC.




(HKRS-2010)

(Leading term QCD expectations)

- HKRS-results at different energies, E_{\perp} and η_{γ} cuts are now available.


t -Acceptance of Roman Pots



Central production with RPs @STAR is taking of. First $pp \rightarrow pXp$ data. Partial wave analysis- soon.

- Phase I set-up focuses on low- t (installed)
- Phase II covers higher- t range (planned to be installed in 2013)

KMR-02



Ongoing **HKRS** studies for **RHIC** energies and kinematics. Correlations between transverse momenta of outgoing protons

Figure 3: The dependence of the survival probability, S^2 , of the rapidity gaps on the azimuthal angle ϕ between the transverse momenta \vec{p}_{it} of the forward going protons in the process $pp \rightarrow p + M + p$, for typical values of p_{1t} and p_{2t} .

Interesting to compare the results for different $\chi_{c0,1,2}$ states.

CENTRAL DIFFRACTION AT THE LHCb

LHCb IS IDEAL FOR DETECTING AND ANALYSING LOW MASS CENTRAL DIFFRACTIVE PRODUCTION OF EXCLUSIVE $\pi^+\pi^-/K^+K^-$ STATES IN:

$$pp \rightarrow p + M + p$$

glueballs, hybrids, heavy quarkonia: χ_c, χ_b
exotic states....

$\pi^+\pi^-/K^+K^-$ STATES AS SPIN-PARITY ANALYZERS.

HOW TO FACILITATE THIS?

Jerry W. Lamsa and Risto Orava

JINST 4:P11019,2009.

LHCb

Excellent particle ID (pion/Kaon separation), vertex and proper time resolution

THE PROPOSED LHCb FSC LAY-OUT

ADD FSCs AT 20 – 100 METERS ON BOTH SIDES OF IP8 – THE FSCs DETECT SHOWERS FROM THE VERY FORWARD PARTICLES.





Figure 1. The layout of LHCb detectors at the LHC Interaction Point (IP8). The proposed Forward Shower Counters (FSCs) are shown as vertical lines (1 to 8). The locations of the dipole (D) and quadrupole (Q) magnet elements are shown as green (dark) and yellow (light) boxes.

Disclaimer : up to the experts to deliver a verdict



Conclusion

- CEP processes observed at the Tevatron can serve as ‘standard candles’ for new physics CEP at the LHC.
- Possibility that χ_{c1} and χ_{c2} CEP may contribute to CDF χ_c events.
- Cannot currently distinguish states, but may be possible with:
 - More detailed analysis and/or higher statistics.
 - Forward proton detection.
 - Different decay modes, $\chi_c \rightarrow \pi\pi, KK, \bar{p}p, \Lambda\bar{\Lambda}$.
- χ_b , dijet, diphton CEP- rich program of studies at the LHC; promising potential of LHCb.
- New STAR@RHIC results on CEP with tagged forward protons soon to come. 
Prospects of CDP studies at ALICE & LHCb

Currently active studies are in progress (both in theory and experiment).



Thank You



BACKUP

UNCERTAINTIES

Known Unknowns

- N(N)LO- radiative effects (K-factors etc..)
 - ‘...possible inadequacy of PT theory in α_s ...’ R.Barbieri et al-1980
 - ‘Right’ choice of gluon densities, in particular at so low scales as in the χ_c case (potentiality of a factor of ~ 3 rise for the H-case).
- Complete model for calculation of enhanced absorption.
- χ_b -experimental widths, decays...



Unknown Unknowns

- Non- pQCD effects in the meson characteristics.
Currently no complete description of heavy quarkonium characteristics.
‘Two gluon width does not tell the whole story.’
- Gluons at so low scales, surprises are not excluded at all.



Factor of 5 up or down
(at best)

	E_{cut}	MRST99	MSTW08LO	M_{min}	MRST99	MSTW08LO
$\sqrt{s} = 7 \text{ TeV}$	5	133	630	10	276	1380
	10	7.32	25.1	20	15.0	53.6
	15	1.15	3.31	30	2.39	7.09
	20	0.274	0.697	40	0.60	1.55
$\sqrt{s} = 10 \text{ TeV}$	5	156	849	10	322	1860
	10	8.77	35.0	20	17.8	74.4
	15	1.43	4.71	30	2.94	10.0
	20	0.34	1.01	40	0.737	2.23
$\sqrt{s} = 14 \text{ TeV}$	5	184	1140	10	378	2470
	10	10.8	48.7	20	21.7	102
	15	1.77	6.71	30	3.59	14.1
	20	0.437	1.47	40	0.934	3.21

Table 8: Central exclusive $\gamma\gamma$ production cross sections (in fb) at different LHC c.m.s energies for different values of cuts on the E_{\perp} ($> E_{\text{cut}}$) of the final-state photons and the invariant mass M_X ($> M_{\text{min}}$) of the diphoton system, in GeV. The photons are restricted to lie in the centre of mass rapidity interval $|\eta_{\gamma}| < 2$.

PROSPECTIVE MEASUREMENTS

- A clear way to resolve the issue of χ_c spin-parity identification will be to search for the two-body decays:

$$Br(\chi_{c0} \rightarrow \pi\pi, K^+K^-) \simeq 1.3\% \quad \chi_{c1}, \eta_c \not\rightarrow \pi\pi, KK \quad Br(\chi_{c2} \rightarrow \pi\pi, K^+K^-) \simeq 0.3\%$$

$$Br(\chi_{c0} \rightarrow p\bar{p}) \simeq 2 * 10^{-4} \quad Br(\chi_{c1} \rightarrow p\bar{p}) \simeq 6.6 * 10^{-5} \quad Br(\chi_{c2} \rightarrow p\bar{p}) \simeq 6.7 * 10^{-5}$$


$$Br(\eta_c \rightarrow p\bar{p}) \simeq 0.13\%$$

- Tagged forward protons: spin-parity ID of old and new heavy meson states, detailed tests of absorption effects
- With sufficient statistics of $\gamma\gamma$ CEP, the measurement of the ratio

$$\sigma(\chi_b) / \sigma(\gamma\gamma)$$
 can be quite instructive (the same mass range, various uncertainties cancel).

Are the early LHC runs, **without** proton taggers, able to check estimates for $pp \rightarrow p+A+p$?

KMR: 0802.0177



Possible checks of:

(i) survival factor S^2 : $W+\text{gaps}, Z+\text{gaps}$

(ii) generalised gluon f_g : $\gamma p \rightarrow Yp$


(iii) Sudakov factor T : 3 central jets

(iv) soft-hard factorisation (enhanced absorptive corrⁿ) $\frac{\#(A+\text{gap}) \text{ evts}}{\#(\text{inclusive } A) \text{ evts}}$
with $A = W, \text{ dijet}, Y, \dots$



Divide et Impera

Exclusive $\gamma\gamma$ Production



Method for excl. $\gamma\gamma$ search is calibrated vs excl e^+e^- analysis:

3 candidates observed:

2 events are good $\gamma\gamma$ candidates

1 event is good $\pi^0\pi^0$ candidate

Theoretical Prediction:


V.A.Khoze et al. Eur. Phys. J C38, 475 (2005)

σ (with our cuts) = (36 + 72 - 24) fb

= 0.8 + 1.6 - 0.5 events.

Cannot yet claim "discovery" as b/g study *a posteriori*,

2 events correspond to $\sigma \sim 90$ fb, agreeing with *Khoze et al.*





$E_T(\gamma) > 5$ GeV

$|\eta(\gamma)| < 1.0$

Exclusive Dimuon Production


Many Physics Processes in this data:



exclusive χ_c in DPE

- Observation of exclusive χ_c PRL 102 242001 (2009)

Exclusive $\chi_c \rightarrow J/\psi (\rightarrow \mu^+ \mu^-) + \gamma$



\rightarrow Allowing EM towers ($E_T > 80$ MeV)
 large increase in the J/ψ peak
 minor change in the $\psi(2s)$ peak



Evidence for

$\chi_c \rightarrow J/\psi + \gamma$ production

$d\sigma/dy|_{y=0} = 75 \pm 14$ nb,
 compatible with theoretical predictions
 160 nb (Yuan 01)
 90 nb (KMR01)

Diffraction with Forward Shower Counters FSC

Mike Albrow, Fermilab

What: We propose to install a set of scintillation counters around both outgoing beam pipes at CMS, $\sim 60\text{m} - 100\text{ m}$


Why:

- ! ? {
- (a) As veto in Level 1 diff. triggers to reduce useless pile-up events
 - (b) To detect rapidity gaps in diffractive events (p or no-p).
 - (c) Measure “low” mass diffraction and double pomeron exchange.
 - (d) Measure σ_{INEL} (if luminosity known, e.g. by Van der Meer)
 - (e) Help establish exclusivity in central exclusive channels
 - (f) To monitor beam conditions on incoming and outgoing beams.
 - (g) To test forward flux simulations (MARS etc.)
 - (h) Additional Luminosity monitor.


Also: They may provide valuable tests of radiation environment to be expected for HPS = High Precision Spectrometers

Central events (0-bias trigger) with forward rap-gaps
 (FSC, ZDC, CASTOR, HF)
 studied for generic Double Pomeron Exchange processes (~ 0.1 mb)


Low mass DPE:



Even without seeing quasi-elastic protons,
 gaps $\gtrsim 4$ units select DPE



The same idea:



Diffraction states with $M > 5$ GeV are very efficiently detected by “OR” of FSC

